



DEPARTMENT OF THE AIR FORCE
59TH MEDICAL WING (AETC)
JOINT BASE SAN ANTONIO - LACKLAND TEXAS

6 FEB 2017

MEMORANDUM FOR SGOZ

ATTN: MAJ BRYANT J. WEBBER

FROM: 59 MDW/SGVU

SUBJECT: Professional Presentation Approval

1. Your paper, entitled **Prevalence and Seroprevalence of Trypanosoma Cruzi Infection in A Military Population in Texas** presented at/published to **AM J Trop Med Hyg (if rejected, then Emerging Infectious Dis, J Infect Dis, PLoS Neglected Trop Dis, or Am J Prev Med)** in accordance with MDWI 41-108, has been approved and assigned local file #**17056**.
2. Pertinent biographic information (name of author(s), title, etc.) has been entered into our computer file. Please advise us (by phone or mail) that your presentation was given. At that time, we will need the date (month, day and year) along with the location of your presentation. It is important to update this information so that we can provide quality support for you, your department, and the Medical Center commander. This information is used to document the scholarly activities of our professional staff and students, which is an essential component of Wilford Hall Ambulatory Surgical Center (WHASC) internship and residency programs.
3. Please know that if you are a Graduate Health Sciences Education student and your department has told you they cannot fund your publication, the 59th Clinical Research Division may pay for your basic journal publishing charges (to include costs for tables and black and white photos). We cannot pay for reprints. If you are 59 MDW staff member, we can forward your request for funds to the designated wing POC.
4. Congratulations, and thank you for your efforts and time. Your contributions are vital to the medical mission. We look forward to assisting you in your future publication/presentation efforts.

Linda Steel-Goodwin

LINDA STEEL-GOODWIN, Col, USAF, BSC
Director, Clinical Investigations & Research Support

PROCESSING OF PROFESSIONAL MEDICAL RESEARCH/TECHNICAL PUBLICATIONS/PRESENTATIONS

INSTRUCTIONS

USE ONLY THE MOST CURRENT 59 MDW FORM 3039 LOCATED ON AF E-PUBLISHING

1. The author must complete page two of this form:
 - a. In Section 2, add the funding source for your study [e.g., 59 MDW CRD Graduate Health Sciences Education (GHSE) (SG5 O&M); SG5 R&D; Tri-Service Nursing Research Program (TSNRP); Defense Medical Research & Development Program (DMRDP); NIH; Congressionally Directed Medical Research Program (CDMRP) ; Grants; etc.]
 - b. In Section 2, there may be funding available for journal costs, if your department is not paying for figures, tables or photographs for your publication. Please state "YES" or "NO" in Section 2 of the form, if you need publication funding support.
2. Print your name, rank/grade, sign and date the form in the author's signature block or use an electronic signature.
3. Attach a copy of the 59 MDW IRB or IACUC approval letter for the research related study. If this is a technical publication/presentation, state the type (e.g. case report, QA/QI study, program evaluation study, informational report/briefing, etc.) in the "Protocol Title" box.
4. Attach a copy of your abstract, paper, poster and other supporting documentation.
5. Save and forward, via email, the processing form and all supporting documentation to your unit commander, program director or immediate supervisor for review/approval.
6. On page 2, have either your unit commander, program director or immediate supervisor:
 - a. Print their name, rank/grade, title; sign and date the form in the approving authority's signature block or use an electronic signature.
7. Submit your completed form and all supporting documentation to the CRD for processing (59crdpubspres@us.af.mil). **This should be accomplished no later than 30 days before final clearance is required to publish/present your materials.** If you have any questions or concerns, please contact the 59 CRD/Publications and Presentations Section at 292-7141 for assistance.
8. The 59 CRD/Publications and Presentations Section will route the request form to clinical investigations, 502 ISG/JAC (Ethics Review) and Public Affairs (59 MDW/PA) for review and then forward you a final letter of approval or disapproval.
9. Once your manuscript, poster or presentation has been approved for a one-time public release, you may proceed with your publication or presentation submission activities, as stated on this form. **Note:** For each new release of medical research or technical information as a publication/presentation, a new 59 MDW Form 3039 must be submitted for review and approval.
10. If your manuscript is accepted for scientific publication, please contact the 59 CRD/Publications and Presentations Section at 292-7141. This information is reported to the 59 MDW/CC. All medical research or technical information publications/presentations must be reported to the Defense Technical Information Center (DITC). See 59 MDWI 41-108, *Presentation and Publication of Medical and Technical Papers*, for additional information.
11. The Joint Ethics Regulation (JER) DoD 5500.07-R, *Standards of Conduct*, provides standards of ethical conduct for all DoD personnel and their interactions with other non-DoD entities, organizations, societies, conferences, etc. Part of the Form 3039 review and approval process includes a legal ethics review to address any potential conflicts related to DoD personnel participating in non-DoD sponsored conferences, professional meetings, publication/presentation disclosures to domestic and foreign audiences, DoD personnel accepting non-DoD contributions, awards, honoraria, gifts, etc. The specific circumstances for your presentation will determine whether a legal review is necessary. **If you (as the author) or your supervisor check "NO" in block 17 of the Form 3039, your research or technical documents will not be forwarded to the 502 ISG/JAC legal office for an ethics review.** To assist you in making this decision about whether to request a legal review, the following examples are provided as a guideline:

For presentations before professional societies and like organizations, the 59 MDW Public Affairs Office (PAO) will provide the needed review to ensure proper disclaimers are included and the subject matter of the presentation does not create any cause for DoD concern.

If the sponsor of a conference or meeting is a DoD entity, an ethics review of your presentation is not required, since the DoD entity is responsible to obtain all approvals for the event.

If the sponsor of a conference or meeting is a non-DoD commercial entity or an entity seeking to do business with the government, then your presentation should have an ethics review.

If your travel is being paid for (in whole or in part) by a non-Federal entity (someone other than the government), a legal ethics review is needed. These requests for legal review should come through the 59 MDW Gifts and Grants Office to 502 ISG/JAC.

If you are receiving an honorarium or payment for speaking, a legal ethics review is required.

If you (as the author) or your supervisor check "YES" in block 17 of the Form 3039, your research or technical documents will be forwarded simultaneously to the 502 ISG/JAC legal office and PAO for review to help reduce turn-around time. If you have any questions regarding legal reviews, please contact the legal office at (210) 671-5795/3365, DSN 473.

NOTE: All abstracts, papers, posters, etc., should contain the following disclaimer statement:

"The views expressed are those of the [author(s)] [presenter(s)] and do not reflect the official views or policy of the Department of Defense or its Components"

NOTE: All abstracts, papers, posters, etc., should contain the following disclaimer statement for research involving humans:

"The voluntary, fully informed consent of the subjects used in this research was obtained as required by 32 CFR 219 and DODI 3216.02_AFI 40-402."

NOTE: All abstracts, papers, posters, etc., should contain the following disclaimer statement for research involving animals, as required by AFMAN 40-401_IP :

"The experiments reported herein were conducted according to the principles set forth in the National Institute of Health Publication No. 80-23, Guide for the Care and Use of Laboratory Animals and the Animal Welfare Act of 1966, as amended."

PROCESSING OF PROFESSIONAL MEDICAL RESEARCH/TECHNICAL PUBLICATIONS/PRESENTATIONS			
1. TO: CLINICAL RESEARCH	2. FROM: (Author's Name, Rank, Grade, Office Symbol) Bryant J. Webber, Maj, USAF, MC, 559 THLS/SGOZ	3. GME/GHSE STUDENT: <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO	4. PROTOCOL NUMBER: FWH20140074H
5. PROTOCOL TITLE: (NOTE: For each new release of medical research or technical information as a publication/presentation, a new 59 MDW Form 3039 must be submitted for review and approval.) The Impact of Chagas Disease on Military Training in Texas			
6. TITLE OF MATERIAL TO BE PUBLISHED OR PRESENTED: Prevalence and Seroprevalence of Trypanosoma cruzi Infection in a Military Population in Texas			
7. FUNDING RECEIVED FOR THIS STUDY? <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO FUNDING SOURCE: AFMSA			
8. DO YOU NEED FUNDING SUPPORT FOR PUBLICATION PURPOSES: <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO			
9. IS THIS MATERIAL CLASSIFIED? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO			
10. IS THIS MATERIAL SUBJECT TO ANY LEGAL RESTRICTIONS FOR PUBLICATION OR PRESENTATION THROUGH A COLLABORATIVE RESEARCH AND DEVELOPMENT AGREEMENT (CRADA), MATERIAL TRANSFER AGREEMENT (MTA), INTELLECTUAL PROPERTY RIGHTS AGREEMENT ETC.? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO NOTE: If the answer is YES then attach a copy of the Agreement to the Publications/Presentations Request Form.			
11. MATERIAL IS FOR: <input checked="" type="checkbox"/> DOMESTIC RELEASE <input type="checkbox"/> FOREIGN RELEASE CHECK APPROPRIATE BOX OR BOXES FOR APPROVAL WITH THIS REQUEST. ATTACH COPY OF MATERIAL TO BE PUBLISHED/PRESENTED.			
<input checked="" type="checkbox"/> 11a. PUBLICATION/JOURNAL (List intended publication/journal.) Am J Trop Med Hyg (if rejected, then Emerging Infectious Dis, J Infect Dis, PLoS Neglected Trop Dis, or Am J Prev Med)			
<input type="checkbox"/> 11b. PUBLISHED ABSTRACT (List intended journal.)			
<input type="checkbox"/> 11c. POSTER (To be demonstrated at meeting: name of meeting, city, state, and date of meeting.)			
<input type="checkbox"/> 11d. PLATFORM PRESENTATION (At civilian institutions: name of meeting, state, and date of meeting.)			
<input type="checkbox"/> 11e. OTHER (Describe: name of meeting, city, state, and date of meeting.)			
12. HAVE YOUR ATTACHED RESEARCH/TECHNICAL MATERIALS BEEN PREVIOUSLY APPROVED TO BE PUBLISHED/PRESENTED? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO ASSIGNED FILE # _____ DATE _____			
13. EXPECTED DATE WHEN YOU WILL NEED THE CRD TO SUBMIT YOUR CLEARED PRESENTATION/PUBLICATION TO DTIC NOTE: All publications/presentations are required to be placed in the Defense Technical Information Center (DTIC).			
DATE March 31, 2017			
14. 59 MDW PRIMARY POINT OF CONTACT (Last Name, First Name, M.I., email) Webber, Bryant J. (bryant.j.webber.mil@mail.mil)			15. DUTY PHONE/PAGER NUMBER 671-4087
16. AUTHORSHIP AND CO-AUTHOR(S) List in the order they will appear in the manuscript.			
LAST NAME, FIRST NAME AND M.I.	GRADE/RANK	SQUADRON/GROUP/OFFICE SYMBOL	INSTITUTION (If not 59 MDW)
a. Primary/Corresponding Author Webber, Bryant J.	Maj	559 THLS/ 559 MDG/ SGOZ	
b. Pawlak, Mary T.	Capt	559 THLS/ 559 MDG/ SGOZ	
c. Valtier, Sandra	GS13	59 MDW/ ST/ CAMD	
d. Daniels, Candelaria C.	MAJ	Department of Clinical Investigation	BAMC
e. Tully, Charla C.	Maj	Department of Infectious Disease	86th Medical Wing
17. IS A 502 ISG/JAC ETHICS REVIEW REQUIRED (JER DOD 5500.07-R)? <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO			
I CERTIFY ANY HUMAN OR ANIMAL RESEARCH RELATED STUDIES WERE APPROVED AND PERFORMED IN STRICT ACCORDANCE WITH 32 CFR 219, AFMAN 40-401_IP, AND 59 MDWI 41-108. I HAVE READ THE FINAL VERSION OF THE ATTACHED MATERIAL AND CERTIFY THAT IT IS AN ACCURATE MANUSCRIPT FOR PUBLICATION AND/OR PRESENTATION.			
18. AUTHOR'S PRINTED NAME, RANK, GRADE Bryant J. Webber, Maj, O-4		19. AUTHOR'S SIGNATURE WEBBER BRYANT J. 1270119383	20. DATE January 25, 2017
21. APPROVING AUTHORITY'S PRINTED NAME, RANK, TITLE Maria J. Belmonte, Lt Col, O-5		22. APPROVING AUTHORITY'S SIGNATURE BELMONTE MARIA J. 1022954527	23. DATE January 25, 2017

PROCESSING OF PROFESSIONAL MEDICAL RESEARCH/TECHNICAL PUBLICATIONS/PRESENTATIONS		
1st ENDORSEMENT (59 MDW/SGVU Use Only)		
TO: Clinical Research Division 59 MDW/CRD Contact 292-7141 for email instructions.	24. DATE RECEIVED January 27, 2017	25. ASSIGNED PROCESSING REQUEST FILE NUMBER 17056
26. DATE REVIEWED January 31, 2017		27. DATE FORWARDED TO 502 ISG/JAC
28. AUTHOR CONTACTED FOR RECOMMENDED OR NECESSARY CHANGES: <input checked="" type="checkbox"/> NO <input type="checkbox"/> YES If yes, give date. _____ <input type="checkbox"/> N/A		
29. COMMENTS <input checked="" type="checkbox"/> APPROVED <input type="checkbox"/> DISAPPROVED IRB approved study with all appropriate disclaimers, and no COI noted. Approved		
30. PRINTED NAME, RANK/GRADE, TITLE OF REVIEWER Kevin Kupferer/GS13/Human Research Subject Protection Expert	31. REVIEWER SIGNATURE KUPFERER,KEVIN R.1086667270 <small>Digitally signed by KUPFERER,KEVIN R.1086667270 DN: cn=US, o=US Government, ou=OSD, ou=PM, ou=USAP, email=KUPFERER,KEVIN R.1086667270 Date: 2017.01.31 08:25:44 -0500</small>	
32. DATE January 31, 2017		
2nd ENDORSEMENT (502 ISG/JAC Use Only)		
33. DATE RECEIVED		34. DATE FORWARDED TO 59 MDW/PA
35. COMMENTS <input type="checkbox"/> APPROVED (In compliance with security and policy review directives.) <input type="checkbox"/> DISAPPROVED		
36. PRINTED NAME, RANK/GRADE, TITLE OF REVIEWER	37. REVIEWER SIGNATURE	
38. DATE		
3rd ENDORSEMENT (59 MDW/PA Use Only)		
39. DATE RECEIVED February 02, 2017		40. DATE FORWARDED TO 59 MDW/SGVU February 03, 2017
41. COMMENTS <input checked="" type="checkbox"/> APPROVED (In compliance with security and policy review directives.) <input type="checkbox"/> DISAPPROVED		
42. PRINTED NAME, RANK/GRADE, TITLE OF REVIEWER Kevin Iinuma, SSgt/E-5, 59 MDW Public Affairs	43. REVIEWER SIGNATURE IINUMA,KEVIN.MITSUGU.1296227 613 <small>Digitally signed by IINUMA,KEVIN.MITSUGU.1296227613 DN: cn=US, o=US Government, ou=OSD, ou=PM, ou=USAP, email=IINUMA,KEVIN.MITSUGU.1296227613 Date: 2017.02.02 07:58:25 -0500</small>	
44. DATE February 03, 2017		
4th ENDORSEMENT (59 MDW/SGVU Use Only)		
45. DATE RECEIVED		46. SENIOR AUTHOR NOTIFIED BY PHONE OF APPROVAL OR DISAPPROVAL <input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> COULD NOT BE REACHED <input type="checkbox"/> LEFT MESSAGE
47. COMMENTS <input type="checkbox"/> APPROVED <input type="checkbox"/> DISAPPROVED		
48. PRINTED NAME, RANK/GRADE, TITLE OF REVIEWER	49. REVIEWER SIGNATURE	
50. DATE		

Prevalence and Seroprevalence of *Trypanosoma cruzi* Infection in a Military Population in Texas

Bryant J. Webber, MD, MPH^{1*}

Mary T. Pawlak, MD, MPH¹

Sandra Valtier, PhD¹

Candelaria C. Daniels, PhD²

Charla C. Tully, DO³

Edward J. Wozniak, DVM, PhD, MPH⁴

Walter D. Roachell, MS⁵

Francisco X. Sanchez Jr., MPH⁵

Audra A. Blasi, DVM¹

Thomas L. Cropper, DVM, MPVM¹

¹59th Medical Wing, Joint Base San Antonio – Lackland, Texas

²Brooke Army Medical Center, Joint Base San Antonio – Fort Sam Houston, Texas

³Landstuhl Regional Medical Center, Kaiserslautern, Germany

⁴Texas State Guard Medical Brigade Headquarters, Camp Mabry, Texas

⁵US Army Public Health Command Central, Joint Base San Antonio – Fort Sam Houston, Texas

*6612 Truemper St. Bldg 6612 Rm 930, Joint Base San Antonio - Lackland, TX 78236;

bryant.j.webber.mil@mail.mil; 210-671-4087

Key Words: *Trypanosoma cruzi*; Chagas disease; triatomine; military training

Abstract Word Count: 142

Text Word Count: 2381

Tables: 1

Abstract

Hundreds of triatomine insects were recently collected at field training sites on Joint Base San Antonio, a large military installation located in south-central Texas. Over 26% of these triatomines tested positive for *Trypanosoma cruzi*, the causative parasitic agent of Chagas disease, and 33% had detectable human blood in their midgut. Given the potential for vector-borne human Chagas disease, a cross-sectional study was conducted to determine the prevalence and seroprevalence of *T. cruzi* infection in highest risk subpopulations on the installation, including students and instructors who work and sleep in triatomine-endemic field settings. Real-time polymerase chain reaction, enzyme-linked immunosorbent assay, and indirect immunofluorescent assay were performed on enrolled subjects (N=1,033), none of whom tested positive for *T. cruzi* or anti-*T. cruzi* antibodies. Current countermeasures employed during field training on Joint Base San Antonio appear to be sufficient for preventing autochthonous human Chagas disease.

Introduction

Chagas disease, or American trypanosomiasis, is caused by infection with *Trypanosoma cruzi*. The protozoan parasite is transmitted to humans most commonly through the infected excreta of hematophagous triatomine insects of the family Reduviidae, entering the bloodstream through a wound or mucous membrane.¹ Known colloquially as “kissing bugs,” triatomines are found throughout the Western hemisphere. Eleven species are endemic to the southern United States, most of which are competent vectors of the parasite.² In addition to the vector-borne route, *T. cruzi* may be transmitted congenitally, orally in contaminated food or beverages, or directly via blood transfusion or organ and tissue transplantation.³

The majority of human cases are subclinical in both the acute and chronic stages, resulting in lifelong, undiagnosed infection. Cardiac disease, gastrointestinal disease, or both develop in approximately one-third of cases, typically manifesting years or decades after initial infection.⁴ Chagas disease mortality is usually attributed to heart failure or ventricular arrhythmia,⁵ but even asymptomatic infection with *T. cruzi* may increase all-cause mortality risk.⁶

Between 5 and 8 million people globally are infected with *T. cruzi*,^{1,7} incurring an annual economic burden of 7.2 billion USD.⁸ Although vectorial transmission is restricted to the Americas,⁹ human migration from endemic to non-endemic countries—and within endemic countries from rural to urban areas—has broadened the distribution of prevalent Chagas disease.¹⁰ Both imported and autochthonous cases occur in the United States, with the former predominating: at least 240 thousand Latin American immigrants are presumably infected,¹¹ whereas fewer than 30 locally-acquired infections have ever been reported.¹² The dearth of documented autochthonous cases, however, may be more indicative of provider unawareness and suboptimal surveillance than true disease incidence.^{12,13} Recognizing the potential for vector-borne transmission across a broad swath of the southern United States, the Centers for Disease Control and Prevention (CDC) prioritizes Chagas as one of five neglected parasitic infections¹⁴ and urges more research to define autochthonous infection risk.¹²

In the greater San Antonio metropolitan area and throughout south-central Texas, Chagas disease has been a known but vaguely defined human disease threat since at least the 1960s.¹⁵⁻¹⁷ An obligation to elucidate that threat recently emerged from several biosurveillance findings on Joint Base San Antonio (JBSA), one of the largest military training installations in the United States. A 2007 serosurvey of military working dogs, all of which are trained on JBSA, found that 24 (8%) harbored anti-*T. cruzi* antibodies. Multiple military working dogs serving in Iraq required evacuation due to cardiomyopathy, later attributed to Chagas disease.¹⁸ A faunal survey was commissioned, which found that 43% (88/205) of collected adult triatomines and 22% (163/736) of nymphs tested positive for *T. cruzi* on polymerase chain reaction (PCR), and blood meal analysis revealed that 33% (43/131) contained human blood in their midgut. Among adults, *Triatoma sanguisuga* (66%) and *Triatoma gerstaeckeri* (30%) were the most common species identified (Daniels, C., unpublished data). This prompted the enforcement of new administrative, technical, and personal protective measures—as well as the reinforcement of existent measures—to protect humans and dogs against vector-borne pathogen exposure during field exercises on JBSA. Since triatomines were often collected in close proximity to high volume training sites, including within field training tents, we also initiated this study to determine human infection risk.

Materials and Methods

This cross-sectional study was designed to establish the prevalence of *T. cruzi* parasitemia and seroprevalence of anti-*T. cruzi* antibodies in five subpopulations most at risk for vector-borne infection while training and working on the installation: students graduating from the US Air Force Security Forces Apprentice course, all of whom had spent 3 weeks training outdoors in a triatomine-endemic area in the month prior to study enrollment, and most of whom had completed a week-long field training exercise on a separate triatomine-endemic site of the installation; instructors from the US Air Force Security Forces Apprentice course; instructors from the US Air Force Basic Military Training field training course; instructors from the Department of Defense Military Working Dog school; and instructors from the US Air Force Survival, Evasion, Resistance, and Escape course. Given reduced prevalence of triatomines during the winter months,¹⁹ and thus reduced likelihood of detecting parasitemia and anti-*T. cruzi* IgM antibodies, we suspended enrollment from December through March.

We administered a questionnaire to all consented participants in order to gather demographic data, quantify exposure risk, and ascertain the geographic location of infection, should a subject test positive. Demographic data included age, sex, and self-reported race and ethnicity. The questionnaire initially focused on vectorial transmission risk by extracting information regarding military training; permanent residence in and travel to triatomine-endemic areas;^{1,2} camping, hunting,²⁰ and exposure to reservoir wildlife^{3,17} in triatomine-endemic areas; and bites by triatomines or by unidentified insects that may have been triatomines. We displayed high-resolution photos of *T. sanguisuga* and *T. gerstaeckeri* adults to facilitate an accurate bite history. After discussing preliminary results with two external consultants, we added questions pertaining to blood transfusion and congenital transmission routes.

We collected whole blood from consented volunteers by peripheral venipuncture. On all subjects from whom we could obtain sufficient aliquots, we performed real-time PCR to determine the prevalence of *T. cruzi* parasitemia and an enzyme-linked immunosorbent assay (ELISA) and indirect immunofluorescent antibody (IFA) test to determine the seroprevalence of anti-*T. cruzi* antibodies.

PCR was conducted per CDC published methodology, utilizing the same primers and probes.²¹ Two multiplex TaqMan assays were performed in parallel targeting three highly conserved and repetitive *T. cruzi* genomic regions: nuclear mini-satellite TCZ; kinetoplast DNA; and the small subunit ribosomal RNA (18S rRNA) gene. An internal validation study was completed using human blood spiked with known amounts of *T. cruzi* epimastigotes. The test was considered positive if all three targets were positive.

The Chagatest ELISA recombinante v.3.0 (Wiener Laboratórios, Rosario, Argentina) was performed, as directed by the manufacturer, for detecting human IgG and IgM anti-*T. cruzi* antibodies. Collected serum samples were incubated with immobilized antigen, washed, incubated with goat anti-human IgG conjugated to horse radish peroxidase, and washed again. Tetramethylbenzidine and hydrogen peroxide were then added, and the reactions were stopped with 2 N sulfuric acid. The colorimetric readings were taken at 450 nm in a plate reader, using a reference wavelength of 650 nm. Cut-off values were defined, per manufacturer guidance, as the mean negative control readings plus 0.3. The test was considered positive if greater than the cut-off value plus 10%, negative if less than the cut-off value minus 10%, and equivocal if within $\pm 10\%$ of the cut-off value.

Sera were also evaluated for the presence of human IgG antibodies against *T. cruzi* via IFA technique. A positive and negative control were validated from the 21-member panel of the SeraCare Life Sciences Chagas Titer AccuSet™. Panel member 1, which had an antibody titer of 1:4096, was used as a positive control at a 1:200 dilution. Panel member 21, which had a negative titer ($<1:128$), was used as a negative control, also at a 1:200 dilution. Subject samples were screened at a 1:128 dilution. Dilutions were placed on *T. cruzi* antigen-coated microscope slides, washed to remove unbound serum antibodies, stained with a fluorescein isothiocyanate-labeled goat anti-human IgG conjugate, and visualized through a fluorescence microscope. A sample was considered positive at a titer equal to or greater than 1:128.

We used descriptive statistics to build demographic and exposure profiles of the enrolled sample, both collectively and stratified by student and instructor status. We compared exposure time between students and instructors with an unpaired t-test, using Epi Info v7.0 (CDC, Atlanta, GA). This study was

approved by the 59th Medical Wing Institutional Review Board (FWH20140074H). Written informed consent was obtained from all subjects.

Results

A total of 1,033 subjects were enrolled. Consistent with the ratio of students to instructors on the installation, the vast majority of subjects (93.1%) were students graduating from the Security Forces Apprentice course (Table 1). During the 16-month study period (April-November 2015 and April-November 2016), we enrolled approximately 15% and 30% of eligible students and instructors, respectively. Most subjects were male (76.9%) and white, non-Hispanic (54.8%). The mean (standard deviation [SD]) age was 21.6 (4.6) years. Three subjects experienced presyncope with venipuncture, all of whom recovered fully without medical intervention. No other adverse events were noted.

Five subjects (0.5%) reported a triatomine bite and 131 (12.7%) reported a bite from an unidentified insect that may have been a triatomine. Subjects experienced 8,130 weeks of total exposure time in the triatomine-endemic field environment of JBSA, for a mean (SD) of 7.7 (18.0) weeks. Instructors (47.0 [45.6] weeks) had a greater mean exposure time than students (4.0 [0.4] weeks) ($p<0.001$). Details on demography, binary risk factors, and time residing and conducting higher risk activities in triatomine-endemic areas are provided in Table 1.

All PCR ($n=1017$), ELISA ($n=1023$), and IFA ($n=1023$) tests were negative, with the exception of one equivocal ELISA result. The enrollment total exceeds laboratory result figures because adequate blood specimens could not be obtained on every subject. The indeterminate ELISA result (0.279 IV [equivocal range: 0.27-0.33 IV]) belonged to a student of Hispanic ethnicity, who was born and lived in Central America for 2 years before emigration to the United States. He then lived in the southwest United States for 19 years prior to arrival at JBSA. During training he experienced three bites that may have been from triatomines, although he could not definitively classify the insect. Repeat ELISA testing was also equivocal, and his PCR and IFA testing were negative. He was advised to visit his health care provider for further discussion and workup.

Discussion

Of 1,033 enrolled subjects, none tested positive for either *T. cruzi* parasites or anti-*T. cruzi* antibodies, suggesting that Chagas disease is currently a negligible threat for military personnel on JBSA. One subject, who tested negative on PCR and IFA, had an equivocal ELISA result. Even if he were truly infected, his case could not be conclusively categorized either as autochthonous, because of his early childhood spent in Latin America, or as vectorial, given the possibility of vertical transmission.

The apparent mismatch between these reassuring findings and the troubling biosurveillance signals—to include cases of canine Chagasic cardiomyopathy and a high volume of *T. cruzi*-infected triatomines—could have several explanations. Exposure of our study subjects to triatomines may have been limited by several antecedent countermeasures: vegetation reduction²² and application of pyrethroid-based insecticides² around tents and field training sites; aggressive reduction of zoonotic reservoirs, particularly woodrats²³ and feral swine;²⁴ requirements pertaining to field uniform wear (i.e., long pants bloused under boots and long sleeves secured at the wrist with buttons); distribution of DEET-based insect repellents free of charge; and, specifically for students during their field training exercise, sleeping in permethrin-treated bed nets.²⁵

These countermeasures alone cannot explain our negative findings, however, since one-third of triatomines collected on our field training sites had fed on humans, and over 13% of our subjects reported bites by triatomines or unidentified insects. Other protective factors may have contributed. First, the stercorarian transmission of parasite from vector to human host is inefficient. Statistical modeling estimates that one case of vectorial transmission requires 900-4,000 contacts with an infected triatomine.²⁶ Second, triatomines opportunistically feed on a variety of vertebrate species—many of which are present on JBSA and have tested positive for *T. cruzi* infection in south-central Texas^{27,28}—offering competitive, non-human blood meal sources. Third, vector competency for any given triatomine species depends on several factors, including environmental distribution, flying and dispersal capacity, inclination to invade human dwellings, and feeding-to-defecation interval.^{2,29,30} Unlike *Triatoma infestans*, *Triatoma dimidiata*, and *Rhodnius prolixus*, the predominant *T. cruzi* vectors in South America,^{2,3} the sylvatic species indigenous to the southern United States are less likely to defecate while taking a blood meal^{31,32} and to

colonize domestic and peridomestic settings.² These biologic and ecologic dynamics may partially explain why autochthonous human Chagas disease appears to be uncommon in the United States,¹² despite a high prevalence of *T. cruzi* infection in triatomine vectors^{18,19,33-35} and substantial evidence for triatomine feeding on humans.^{33,36,37} This incongruity between a high prevalence of triatomines and low seroprevalence of human *T. cruzi* infection was also reported on the Yucatan Peninsula of Mexico.³⁸

Another explanation, albeit a study limitation and not a causative factor, is the possibility of false negative testing. According to a recent meta-analysis, the Weiner ELISA has a sensitivity of 93.7% (95% confidence interval: 87.7%, 96.9%) for detecting human antibodies against *T. cruzi*.³⁹ IFA testing is approximately 90% sensitive, although substantial heterogeneity exists across studies.⁴⁰ This may be due to methodological differences, particularly with titer dilution cutoffs used to determine positive results. Our 1:128 dilution screening was designed to maximize overall test accuracy, but other laboratories may demarcate a positive result at a titer of 1:80,⁴¹ thus conceding some specificity for improved sensitivity. PCR, which is most valuable diagnostically during the parasitemic acute stage of disease,⁴ has a sensitivity below 50% for detecting chronic infection in adults.³⁹ However, by utilizing 3 tests in parallel and analyzing over a thousand subjects, imperfect sensitivity for any one test does not undermine conclusions drawn this study.

The lack of molecular and serologic evidence of *T. cruzi* infection in our study sample, though encouraging, should not be misapplied. First, our findings should not be used to exclude Chagas disease from the differential diagnosis list when evaluating service members who trained or worked on triatomine-endemic field sites of JBSA. Although we selected populations with the highest risk of exposure, we only tested a sample thereof, and our results may not be generalizable to populations on the installation before aggressive countermeasures were deployed. Second, our findings should not be used to establish countermeasure success. In the absence of a control group not employing these measures, our study was not designed to verify their effectiveness. Current countermeasures appear sufficient but not categorically necessary in preventing autochthonous human Chagas disease on JBSA.

Despite an abundance of *T. cruzi*-infected triatomine vectors, some of which evidently feed on humans, Chagas disease is currently not a major infectious disease threat for military students and instructors on JBSA. Even if parasite transmissibility is intrinsically unlikely due to bioecological factors, primary preventive measures reducing exposure to *T. cruzi* and other vector-borne pathogens should continue. In order to stage realistic military field training exercises while maximizing the health of humans, animals, and the environment, we urge holistic One Health approaches built on collaboration between military training leadership, civil engineers, and medical, veterinary, and public health personnel.

Acknowledgements

We thank La'Quita Armstrong-Spenrath, Manuel Caballero, Arlinda Haliti, Sallie Hall, David Hill, Dion Holmes, Samantha Hune, Constance Kowat, Richard McIntosh, Kimberly Murphy, Lois Robinson, Michele Tavish, Juste Tchandja, and John Yevick at the 59th Medical Wing; Bernardo Delgado Jr. and Angel Osuna at the US Army Public Health Command; Susan Montgomery and Theresa Benedict at the CDC; and Melissa Garcia at Baylor College of Medicine.

Financial Support

This study was funded by the US Air Force Medical Services Agency (Falls Church, VA, USA). The authors have no financial conflicts of interest.

Disclosures

The voluntary, fully informed consent of subjects used in this research was obtained as required by 32 CFR 219 and DoDI 3216.02_AFI 40-402. The views expressed are those of the authors and do not reflect the official views or policy of the Department of Defense and its Components. Preliminary results from this study were presented at ID Week; October 7-11, 2015, San Diego, CA; and at the American Society of Tropical Medicine and Hygiene Annual Meeting; November 13-17, 2016, Atlanta, GA.

The views expressed are those of the author(s)/presenter(s) and do not reflect the official view or policy of the Department of Defense or its Components.

Authors

Bryant J. Webber, MD, MPH

233 Trainee Health Surveillance

234 59th Medical Wing

235 Joint Base San Antonio - Lackland, Texas, USA

236 bryant.j.webber.mil@mail.mil

237 *Mary T. Pawlak, MD, MPH*

238 Trainee Health Surveillance

239 59th Medical Wing

240 Joint Base San Antonio - Lackland, Texas, USA

241 mary.t.pawlak.mil@mail.mil

242 *Sandra Valtier, PhD*

243 Center for Advanced Molecular Detection

244 59th Medical Wing

245 Joint Base San Antonio - Lackland, Texas, USA

246 sandra.valtier.civ@mail.mil

247 *Candelaria C. Daniels, PhD*

248 Department of Clinical Investigation

249 Brooke Army Medical Center

250 Joint Base San Antonio - Fort Sam Houston, Texas, USA

251 candelaria.c.daniels.civ@mail.mil

252 *Charla C. Tully, DO*

253 Department of Infectious Disease

254 Landstuhl Regional Medical Center

255 Kaiserslautern, Germany

256 charla.c.tully.mil@mail.mil

257 *Edward J. Wozniak, DVM, PhD, MPH*

258 Texas State Guard Medical Brigade Headquarters

259 Camp Mabry, Texas, USA
 260 edward.wozniak@txsg.state.tx.us
 261 *Walter D. Roachell, MS*
 262 Department of Entomology
 263 US Army Public Health Command Central
 264 Joint Base San Antonio - Fort Sam Houston, Texas, USA
 265 walter.d.roachell.civ@mail.mil
 266 *Francisco X. Sanchez Jr., MPH*
 267 Department of Defense Food Analysis & Diagnostic Laboratory
 268 US Army Public Health Command Central
 269 Joint Base San Antonio - Fort Sam Houston, Texas, USA
 270 francisco.x.sanchez2.civ@mail.mil
 271 *Audra A. Blasi, DVM*
 272 Public Health Flight
 273 59th Medical Wing
 274 Joint Base San Antonio - Lackland, Texas, USA
 275 audra.a.blasi.mil@mail.mil
 276 *Thomas L. Cropper, DVM, MPVM*
 277 Trainee Health Surveillance
 278 59th Medical Wing
 279 Joint Base San Antonio - Lackland, Texas, USA
 280 thomas.l.cropper.civ@mail.mil

281 **References**

- 282 1. Rassi A Jr, Rassi A, Marin-Neto JA, 2010. Chagas disease. *Lancet* 375: 1388-1402.
- 283 2. Klotz SA, Dorn PL, Mosbacher M, Schmidt JO, 2014. Kissing bugs in the United States: risk for
- 284 vector-borne disease in humans. *Environ Health Insights* 10: 49-59.

3. Bern C, Kjos S, Yabsley MJ, Montgomery SP, 2011. *Trypanosoma cruzi* and Chagas' disease in the United States. *Clin Microbiol Rev* 24: 655-681.
4. Bern C, 2015. Chagas' disease. *N Engl J Med* 373: 456-466.
5. Malik LH, Singh GD, Amsterdam EA, 2015. The epidemiology, clinical manifestations, and management of Chagas heart disease. *Clin Cardiol* 38: 565-569.
6. Cucunubá ZM, Okuwoga O, Basáñez MG, Nouvellet P, 2016. Increased mortality attributed to Chagas disease: a systematic review and meta-analysis. *Parasit Vectors* 9: 42.
7. World Health Organization, 2015. Chagas disease in Latin America: an epidemiological update based on 2010 estimates. *Wkly Epidemiol Rec* 90: 33-43.
8. Lee BY, Bacon KM, Bottazzi ME, Hotez PJ, 2013. Global economic burden of Chagas disease: a computational simulation model. *Lancet Infect Dis* 13: 342-348.
9. Liu Q, Zhou XN, 2015. Preventing the transmission of American trypanosomiasis and its spread into non-endemic countries. *Infect Dis Poverty* 28: 60.
10. Conners EE, Vinetz JM, Weeks JR, Brouwer KC, 2016. A global systematic review of Chagas disease prevalence among migrants. *Acta Trop* 156: 68-78.
11. Manne-Goehler J, Umeh CA, Montgomery SP, Wirtz VJ, 2016. Estimating the burden of Chagas disease in the United States. *PLoS Negl Trop Dis* 10: e0005033.
12. Montgomery SP, Parise ME, Dotson EM, Bialek SR, 2016. What Do We Know About Chagas Disease in the United States? *Am J Trop Med Hyg* 95: 1225-1227.
13. Basile L, Jansa JM, Carlier Y, Salamanca DD, Angheben A, Bartoloni A, Seixas J, Van Gool T, Cañavate C, Florez-Chávez M, Jackson Y, Chiodini PL, Albajar-Viñas P; Working Group on Chagas Disease, 2011. Chagas disease in European countries: the challenge of a surveillance system. *Euro Surveill* 16.
14. Montgomery SP, Starr MC, Cantey PT, Edwards MS, Meymandi SK, 2014. Neglected parasitic infections in the United States: Chagas disease. *Am J Trop Med Hyg* 90: 814-818.

15. Lathrop GD, Ominsky AJ, 1965. Chagas' disease study in a group of individuals bitten by North American triatomids. *Aeromed Rev* 9: 1-5.
16. Pippin WF, 1970. The biology and vector capability of *Triatoma sanguisuga texana* usinger and *Triatoma gerstaeckeri* (Stål) compared with *Rhodnius prolixus* (Stål) (Hemiptera: Triatominae). *J Med Entomol* 7: 30-45.
17. Sarkar S, Strutz SE, Frank DM, Rivaldi CL, Sissel B, Sánchez-Cordero V, 2010. Chagas' disease risk in Texas. *PLoS Negl Trop Dis* 4: e836.
18. McPhatter L, Roachell W, Mahmood F, Hoffman L, Lockwood N, Osuna A, Lopez J, Debboun M, 2012. Vector surveillance to determine species composition and occurrence of *trypanosoma cruzi* at three military installations in San Antonio, Texas. *US Army Med Dep J Jul-Sep*: 12-21.
19. Wozniak EJ, Lawrence G, Gorchakov R, Alamgir H, Dotson E, Sissel B, Sarkar S, Murray KO, 2015. The biology of the triatomine bugs native to south central Texas and assessment of the risk they pose for autochthonous Chagas disease exposure. *J Parasitol* 101: 520-528.
20. Garcia MN, Hotez PJ, Murray KO, 2014. Potential novel risk factors for autochthonous and sylvatic transmission of human Chagas disease in the United States. *Parasit Vectors* 7: 311.
21. Qvarnstrom Y, Schijman AG, Veron V, Aznar C, Steurer F, da Silva AJ, 2012. Sensitive and specific detection of *Trypanosoma cruzi* DNA in clinical specimens using a multi-target real-time PCR approach. *PLoS Negl Trop Dis* 6: e1689.
22. Shender L, Niemela M, Conrad P, Goldstein T, Mazet J, 2016. Habitat management to reduce human exposure to *Trypanosoma cruzi* and western conenose bugs (*Triatoma protracta*). *Ecohealth* 13: 525-534.
23. Charles RA, Kjos S, Ellis AE, Barnes JC, Yabsley MJ, 2013. Southern plains woodrats (*Neotoma micropus*) from southern Texas are important reservoirs of two genotypes of *Trypanosoma cruzi* and host of a putative novel *Trypanosoma* species. *Vector Borne Zoonotic Dis* 13: 22-30.

24. Comeaux JM, Curtis-Robles R, Lewis BC, Cummings KJ, Mesenbrink BT, Leland BR, Bodenchuk MJ, Hamer SA, 2016. Survey of Feral Swine (*Sus scrofa*) Infection with the Agent of Chagas Disease (*Trypanosoma cruzi*) in Texas, 2013-2014. *J Wildl Dis* 52: 627-630.
25. Armed Forces Pest Management Board, 2015. Technical Guide No. 36: Personal protective measures against insects and other arthropods of military significance. 6 November 2015. Available at <http://www.acq.osd.mil/eie/afpmb/docs/techguides/tg36.pdf> (accessed January 10, 2017).
26. Nouvellet P, Dumonteil E, Gourbière S, 2013. The improbable transmission of *Trypanosoma cruzi* to human: The missing link in the dynamics and control of Chagas disease. *PLoS Negl Trop Dis* 7: e2505.
27. Curtis-Robles R, Lewis BC, Hamer SA, 2016. High *Trypanosoma cruzi* infection prevalence associated with minimal cardiac pathology among wild carnivores in central Texas. *Int J Parasitol Parasites Wildl* 5: 117-123.
28. Garcia MN, O'Day S, Fisher-Hoch S, Gorchakov R, Patino R, Feria Arroyo TP, Laing ST, Lopez JE, Ingber A, Jones KM, Murray KO, 2016. One Health Interactions of Chagas Disease Vectors, Canid Hosts, and Human Residents along the Texas-Mexico Border. *PLoS Negl Trop Dis* 10: e0005074.
29. Kjos SA, Snowden KF, Olson JK, 2009. Biogeography and *Trypanosoma cruzi* infection prevalence of Chagas disease vectors in Texas, USA. *Vector Borne Zoonotic Dis* 9: 41-50.
30. Martínez-Ibarra JA, Alejandre-Aguilar R, Paredes-González E, Martínez-Silva MA, Solorio-Cibrián M, Noqueda-Torres B, Trujillo-Contreras F, Novelo-López M, 2007. Biology of three species of North American Triatominae (Hemiptera: Reduviidae: Triatominae) fed on rabbits. *Mem Inst Oswaldo Cruz* 102: 925-930.
31. Klotz SA, Doran PL, Klotz JH, Pinnas JL, Weinrach C, Kurtz JR, Schmidt J, 2009. Feeding behavior of triatomines from the southwestern United States: an update on potential risk for transmission of Chagas disease. *Acta Trop* 111: 114-118.
32. Zeledón R, Alvarado R, Jirón LF, 1977. Observations on the feeding and defecation patterns of three triatomine species (Hemiptera: Reduviidae). *Acta Trop* 34: 65-77.

33. Waleckx E, Suarez J, Richards B, Dorn PL, 2014. *Triatoma sanguisuga* blood meals and potential for Chagas disease, Louisiana, USA. *Emerg Infect Dis* 20: 2141-2143.
34. Curtis-Robles R, Wozniak EJ, Auckland LD, Hamer GL, Hamer SA, 2015. Combining public health education and disease ecology research: Using citizen science to assess Chagas disease entomological risk in Texas. *PLoS Negl Trop Dis* 9: e0004235.
35. Buhaya MH, Galvan S, Maldonado RA, 2015. Incidence of *Trypanosoma cruzi* infection in triatomines collected at Indio Mountains Research Station. *Acta Trop* 150: 97-99.
36. Klotz SA, Schmidt JO, Dorn PL, Ivanyi C, Sullivan KR, Stevens L, 2014. Free-roaming kissing bugs, vectors of Chagas disease, feed often on humans in the Southwest. *Am J Med* 127: 421-426.
37. Gorchakov R, Trosclair LP, Wozniak EJ, Fera PT, Garcia MN, Gunter SM, Murray KO, 2016. *Trypanosoma cruzi* infection prevalence and bloodmeal analysis in triatomine vectors of Chagas disease from rural peridomestic locations in Texas, 2013-2014. *J Med Entomol* 53: 911-918.
38. Monteón V, Solis-Oviedo R, Lopez R, Hernández O, Tellez CA, 2015. Low seroprevalence of *Trypanosoma cruzi* infection and chronic chagasic cardiomyopathy in a region with abundance of triatomine vectors in Yucatan Peninsula of Mexico. *Ann Parasitol* 61: 263-267.
39. Brasil PE, Castro R, Castro L, 2016. Commercial enzyme-linked immunosorbent assay versus polymerase chain reaction for the diagnosis of chronic Chagas disease: a systematic review and meta-analysis. *Mem Inst Oswaldo Cruz* 111: 1-19.
40. Afonso AM, Ebell MH, Tarleton RL, 2012. A systematic review of high quality diagnostic tests for Chagas disease. *PLoS Negl Trop Dis* 6: e1881.
41. Umezawa ES, Bastos SF, Coura JR, Levin MJ, Gonzalez A, Rangel-Aldao R, Zingales B, Luquetti AO, da Silveira JF, 2003. An improved serodiagnostic test for Chagas' disease employing a mixture of *Trypanosoma cruzi* recombinant antigens. *Transfusion* 43: 91-97.

383 **Table 1. Demographic and risk factor profile of study subjects (N=1,033)**

	Students N=962	Instructors N=71	Total N=1,033
Population			
-Security Forces Apprentice course students	962 (100%)		962 (93.1%)
-Security Forces Apprentice course instructors		2 (2.8%)	2 (0.2%)
-Basic Military Training field training instructors		36 (50.7%)	36 (3.5%)
-Military Working Dog School instructors		23 (32.4%)	23 (2.2%)
-Survival, Evasion, Resistance, and Escape instructors		10 (14.1%)	10 (1.0%)
Age, mean (std dev)	20.9 (3.6)	31.6 (5.0)	21.6 (4.6)
Sex			
-Male	735 (76.4%)	59 (83.1%)	794 (76.9%)
-Female	227 (23.6%)	12 (16.9%)	239 (23.1%)
Race/ethnicity			
-White, non-Hispanic	515 (53.5%)	51 (71.8%)	566 (54.8%)
-Black, non-Hispanic	121 (12.6%)	7 (9.9%)	128 (12.4%)
-Hispanic	221 (23.0%)	8 (11.3%)	229 (22.2%)
-Other	105 (10.9%)	5 (7.0%)	110 (10.6%)
Potential exposures			
-Known triatomine bite	4 (0.4%)	1 (1.4%)	5 (0.5%)
-Unidentified insect bite	102 (10.6%)	29 (40.8%)	131 (12.7%)
-Received blood products in US*	7 (0.8%)	3 (5.9%)	10 (1.1%)
-Received blood products outside US*	3 (0.3%)	0	3 (0.3%)
-Mother lived in Latin America before birth*	90 (10.2%)	4 (7.8%)	94 (10.1%)
Weeks in triatomine-endemic area, mean (std dev)			
-Field environment at JBSA-Lackland	4.0 (0.4)	47.0 (45.6)	7.7 (18.0)
-Camping/hunting in Latin America or southwest US†	30.8 (125.0)	75.8 (283.5)	35.2 (147.0)
-Wildlife exposure‡ in Latin America or southwest US†	110.2 (280.9)	246.1 (513.7)	126.7 (320.2)
-Living/traveling in Latin America	78.0 (230.1)	80.9 (225.2)	67.3 (211.6)
-Living/traveling in southwest US†	430.4 (485.5)	238.9 (379.4)	380.9 (474.4)

*N=931 since these questions were added to the questionnaire after study initiation.

†Southwest United States was defined as Arizona, California, Colorado, Nevada, New Mexico, Oklahoma, Texas, and Utah.

‡Wildlife exposure was defined as either hunting or living in a dwelling infested by woodrats, raccoons, opossums, skunks, wild hogs, coyotes, or deer.